

What is most important to remember?

- Chlorthalidone is a “water pill” used to treat high blood pressure and remove extra fluid in the body
- It is important to continue taking this medicine even if you feel normal. Most people with high blood pressure do not feel sick
- Do not start any new medicines, over-the-counter drugs or herbal remedies without talking to your doctor
- Contact your doctor immediately if you experience itching, rash, or swelling of the face, lips, tongue or throat, or if you experience very bad dizziness or passing out

If you have questions:

Ask your doctor, nurse or pharmacist for more information about chlorthalidone

Strong Internal Medicine



Strong Internal Medicine
601 Elmwood Avenue
Ambulatory Care Facility, 5th Floor
Rochester, NY 14642
Phone: (585) 275 7424

Visit our website at:
[www.urmc.rochester.edu/medicine/
general-medicine/patientcare/](http://www.urmc.rochester.edu/medicine/general-medicine/patientcare/)

Chlorthalidone: Important Patient Information



What does chlorthalidone do?

- It is “water pill” used to treat high blood pressure and remove extra fluid in the body

How should chlorthalidone be used?

- Take this medicine as directed by your doctor
- It is usually taken one (1) time a day after a meal, preferably after breakfast
- It is best to take this medicine in the morning to avoid going to the bathroom during the night

Who should know I am taking chlorthalidone?

- Keep a list of all your medicines (prescription, natural products, supplements, vitamins, over-the-counter) and give it to your healthcare provider (doctor, nurse, nurse practitioner, pharmacist, or physician assistant)

Are there any interactions with other drugs that I need to worry about?

- There are many drug interactions that may increase your risk of side effects
- Do not start new prescriptions, over-the-counter medicines, or herbal and dietary supplements without telling your doctor

What should I do if I miss a dose of chlorthalidone?

- Take your dose as soon as you remember
- If it is almost time for your next dose, skip the missed dose and go back to your normal dose time
- Do not take 2 doses at the same time or extra doses to make up for a missed dose

What side effects could occur with chlorthalidone?

- Feeling dizzy
- Dry mouth, upset stomach, or throwing up
- Signs of low potassium such as: muscle pain or weakness, muscle cramps, or a heartbeat that does not feel normal

What are some things that I need to be aware of when taking chlorthalidone?

- Before taking this medicine tell your doctor or pharmacist if you have a “sulfa” (sulfonamide) allergy, are allergic to chlorthalidone, or if you are allergic to any other medicines, foods, or substances
- Before using this medicine tell your doctor or pharmacist your medical history, especially if you have diabetes, kidney disease, or liver disease
- Before taking this medicine tell your doctor if you are pregnant, plan to get pregnant, or are breastfeeding. You will need to talk about the benefits and risks of using this medicine
- Chlorthalidone will make you pass urine more often. You should not take it after 6pm to avoid waking up at night to use the bathroom
- Taking this medicine will require routine blood work. Talk with your doctor
- You should have your blood pressure checked often
- Avoid drinking alcohol (includes wine, beer, and liquor) when taking this medicine
- Stand up slowly when getting up to avoid getting dizzy or passing out
- You may get sunburned more easily. Avoid sun, sunlamps, and tanning beds. Use sunscreen and wear clothing and eyewear that protect you from the sun
- It is important to continue taking this medicine even if you feel normal. Most people with high blood pressure do not feel sick
- Store in a cool, dry place. Protect from light and do not store in the bathroom
- Keep out of the reach of children and pets

If you have any of the following side effects, call your doctor or visit the emergency room immediately:

- Signs of low potassium
- Very bad dizziness or passing out
- Very upset stomach or throwing up
- Not able to pass urine
- Any sign of a severe allergic reaction including: rash; hives; red, swollen, blistered or peeling skin with or without fever; wheezing; tightness of chest or throat; trouble breathing or talking; swelling of face, lips, tongue, or throat