

PicoLab® Mouse Diet 20

5058*

DESCRIPTION

PicoLab® Mouse Diet 20 is a Constant Nutrition® formulation providing 20% protein for mouse colonies that require extra levels of energy needed for maximum production in post-partum breeding. Irradiation treatment and special 4-ply packaging provide virtually bacteria-free dietary control.

Features and Benefits

- Formulated with 20% protein for mouse breeding colonies
- Precision processing and selection of highest quality ingredients assures Constant Nutrition® quality
- Designed to meet the energy needs of breeding mouse colonies, transgenic strains, and mice exposed to higher stress levels
- Irradiation gives reliable microbial control and eliminates the need for autoclaving

Product Forms Available

- Oval pellet, 10 mm x 16 mm x 25 mm length (3/8"x5/8"x1")
- Meal (ground pellets), special order

Other Versions Available

- 5062 Pico-Vac® Mouse Diet 20

GUARANTEED ANALYSIS

| | | |
|------------------------------|-------|-------|
| Crude protein not less than | | 20.0% |
| Crude fat not less than | | 9.0% |
| Crude fiber not more than | | 4.0% |
| Ash not more than | | 6.5% |
| Added minerals not more than | | 2.5% |

INGREDIENTS

Ground wheat, ground corn, dehulled soybean meal, wheat germ, fish meal, brewers dried yeast, corn gluten meal, porcine animal fat preserved with BHA, soybean oil, calcium carbonate, salt, dicalcium phosphate, monocalcium phosphate, choline chloride, menadione dimethylpyrimidinol bisulfite, DL-methionine, vitamin A acetate, cholecalciferol, pyridoxine hydrochloride, dried whey, folic acid, dl-alpha tocopheryl acetate, biotin, thiamin mononitrate, calcium pantothenate, lecithin, riboflavin, nicotinic acid, casein, vitamin B₁₂ supplement, manganous oxide, zinc oxide, ferrous carbonate, copper sulfate, zinc sulfate, calcium iodate, cobalt carbonate, sodium selenite.

FEEDING DIRECTIONS

Feed ad libitum to mice. Plenty of fresh, clean water should be available to the animals at all times.

Mice-Adult mice will eat up to 5 grams of pelleted ration daily. Some of the larger strains may eat as much as 8 grams per day per animal. Feed should be available on a free choice basis in wire feeders above the floor of the cage.

CHEMICAL COMPOSITION¹

| | | |
|--|-------|------|
| Nutrients² | | |
| Protein, % | | 21.8 |
| Arginine, % | | 1.15 |
| Cystine, % | | 0.31 |
| Glycine, % | | 0.93 |
| Histidine, % | | 0.50 |
| Isoleucine, % | | 1.02 |
| Leucine, % | | 1.82 |
| Lysine, % | | 1.13 |
| Methionine, % | | 0.67 |
| Phenylalanine, % | | 0.97 |
| Tyrosine, % | | 0.64 |
| Threonine, % | | 0.79 |
| Tryptophan, % | | 0.25 |
| Valine, % | | 1.03 |
| Serine, % | | 1.07 |
| Aspartic Acid, % | | 2.13 |
| Glutamic Acid, % | | 4.47 |
| Alanine, % | | 1.34 |
| Proline, % | | 1.54 |
| Taurine, % | | 0.02 |
| Fat (ether extract), % | | 9.0 |
| Fat (acid hydrolysis), % | | 9.1 |
| Cholesterol, ppm | | 200 |
| Linoleic Acid, % | | 2.32 |
| Linolenic Acid, % | | 0.21 |
| Arachidonic Acid, % | | 0.02 |
| Omega-3 Fatty Acids, % | | 0.32 |
| Total Saturated Fatty Acids, % | | 2.72 |
| Total Monounsaturated Fatty Acids, % | | 2.88 |
| Fiber (Crude), % | | 2.2 |
| Neutral Detergent Fiber ³ , % | | 10.8 |
| Acid Detergent Fiber ⁴ , % | | 3.0 |
| Nitrogen-Free Extract (by difference), % | | 51.8 |
| Starch, % | | 39.3 |
| Glucose, % | | 0.16 |
| Fructose, % | | 0.16 |
| Sucrose, % | | 0.71 |
| Lactose, % | | 0.78 |
| Total Digestible Nutrients, % | | 85.3 |
| Gross Energy, kcal/gm | | 4.60 |
| Physiological Fuel Value⁵, kcal/gm | | 3.75 |
| Metabolizable Energy, kcal/gm | | 3.56 |

Minerals

| | | |
|-----------------------------|-------|------|
| Ash, % | | 5.0 |
| Calcium, % | | 0.81 |
| Phosphorus, % | | 0.60 |
| Phosphorus (non-phytate), % | | 0.33 |
| Potassium, % | | 0.70 |
| Magnesium, % | | 0.16 |

Vitamins

| | | |
|---------------------------------------|-------|-------|
| Carotene, ppm | | Trace |
| Vitamin K (as menadione), ppm | | 3.1 |
| Thiamin Hydrochloride, ppm | | 15 |
| Riboflavin, ppm | | 8.0 |
| Niacin, ppm | | 90 |
| Pantothenic Acid, ppm | | 21 |
| Choline Chloride, ppm | | 2200 |
| Folic Acid, ppm | | 2.9 |
| Pyridoxine, ppm | | 9.6 |
| Biotin, ppm | | 0.30 |
| B ₁₂ , mcg/kg | | 51 |
| Vitamin A, IU/gm | | 15 |
| Vitamin D ₃ (added), IU/gm | | 3.3 |
| Vitamin E, IU/kg | | 57 |
| Ascorbic Acid, mg/gm | | — |

Calories provided by:

| | | |
|------------------------|-------|--------|
| Protein, % | | 23.189 |
| Fat (ether extract), % | | 21.635 |
| Carbohydrates, % | | 55.176 |

*Product Code

1. Based on the latest ingredient analysis information. Since nutrient composition of natural ingredients varies, analysis will differ accordingly.
2. Nutrients expressed as percent of ration except where otherwise indicated. Moisture content is assumed to be 10.0% for the purpose of calculations.
3. NDF = approximately cellulose, hemi-cellulose and lignin.
4. ADF = approximately cellulose and lignin.
5. Physiological Fuel Value (kcal/gm) = Sum of decimal fractions of protein, fat and carbohydrate (use Nitrogen Free Extract) x 4.94 kcal/gm respectively.