

# HENDRICH II FALL RISK MODEL

Form # 304 (8/16)

Name: _____	DOB: _____	Date: _____
Risk Factor		Risk Points
Confusion / disorientation / impulsivity		4
Symptomatic Depression		2
Altered Elimination		1
Dizziness / Vertigo		1
Gender (Male)		1
Any Administered Antiepileptics (anticonvulsants): <small>(Cardamazepine, Divolproex Sodium, Ethotoin, Ethosuximide, Felbamate, Fosphenytoin, Gabapentin, Lamotrigine, Mephenytoin, Methsuximide, Phenobarbital, Phenytoin, Primidone, Topiramate, Trimethadione, Valproic Acid)</small>		2
Any Administered Benzodiazepines: <small>(Alprazolam, Chlordiazepoxide, Clonazepam, Clorazepate Dipotassium, Diazepam, Flurazepam, Halazepam, Lorazepam, Midazolam, Oxazepam, Triazolam)</small>		1
<b>Get Up and Go Test: "Rising from a Chair"</b>		
<b>* If unable to assess, monitor for change in activity level, assess other risk factors, document both on patient chart with date and time.</b>		
Able to rise in a single movement – No loss of balance with steps		0
Pushes up, successful in one attempt		1
Multiple attempts, but successful		3
Unable to rise without assistance during test (OR if a medical order states the same and/or complete bed rest is ordered) *If unable to assess, document this on the patient chart with the date and time.		4
<b>A score of 5 or greater = High risk</b>		<b>Total Score</b>
Nurses Signature ►	Date:	Time:

## HENDRICH II FALL RISK MODEL