

# Confessions of Physicians: What Systemic Reporting Does Not Uncover

Suzanne B. Karan, MD

Jeffrey S. Berger, MD, MBA

Michael Wajda, MD

**21** **I**t's July, and the newest group of anesthesiology residents is sitting around the conference table. With clear memories of previous failed efforts to get them to "open up," the program director is trying something new. Sixteen pieces of folded paper are arranged in a pile, one for each resident in the room. Each paper contains a typed statement, and there are no names attached to these "confessions."

The papers are shuffled and dealt, poker-style, to the group. Participants are reminded, "Even if you get your own statement, do not reveal the authorship, as anonymity is critical to the session."

The first statement is read:

*Been really repulsed by the whole room "turnover" process. Reusing blood pressure cuffs that have questionable stains on them; the EKG wires go straight from the sticky, bloody floor to being hung on top of IV pole for next patient use. What a gross-out this has been . . . yuck.*

The facilitator decides to remain silent for a few moments. After an uncomfortable pause, a resident wonders aloud, "Who cleans the equipment in between cases? How do we know when we have to do it, or when the technician is responsible?" Another resident responds confidently, "I just do it. I mean, my Mom could be the next patient . . . why take any chances?" The conversation lulls and the next resident reads:

*I've had the privilege of being with some pretty serious residents the last couple of weeks. Their actions remind me of stealthy ninjas. I confess there are times when I'm sitting there with monitors alarming off the hilt and have been nudged by the residents who say, "Did you notice that alarm blaring in your ear?" No . . . sadly, no.*

A conversation regarding situational awareness and vigilance ensues. With each confessional statement,

the tension that marked the outset of the session dissipates further. The next confession is read:

*I had 2 instances where I went to get my drug, only to find that I forgot to label the syringe. In both instances, I knew what the drugs were because I had drawn them up myself recently and placed them in their usual places and all the other drugs I had were labeled. I used the drugs, but now looking back, this is probably bad practice and where mistakes happen. Now I make it a habit to label the syringes first before I draw up the drugs.*

What this new approach to resident conversations is trying to achieve is to overcome the stigma of reporting and discussing incidents and situations that negatively affect the quality and safety of care. The most important barrier to improving safety is the lack of a standardized metric of adverse events and incident reports. A few programs have been successful at increasing reporting in certain disciplines<sup>1</sup> and globally,<sup>2</sup> although more is clearly needed.<sup>3</sup> Since the overall incidence of poor-quality outcomes is low, the use of such metrics as report cards and insurance claims to benchmark specific concerns tends to be underpowered. Consequently, there is a general dearth of data that could bring about systematic improvements in patient safety.

**22** The Institute of Medicine drastically "changed the conversation"<sup>3</sup> regarding error reporting by shifting the focus to systems rather than on individuals. In doing so, the goal is to more readily engage all stakeholders, including health care workers, to become part of the solution. Hospital staff is constantly encouraged to report errors and near-misses in event-reporting systems. Although US health care institutions have begun promoting transparency to improve the safety of care, there is no widespread achievement of this goal. Current hospital-based reporting systems are profoundly underused by physicians compared with nurses and other staff<sup>4</sup> (1.1% versus 45.3% and 53.6%, respectively). A perceived threat to autonomy and fear of reprisal

DOI: <http://dx.doi.org/10.4300/JGME-D-15-00054.1>

## BOX RULES FOR CONFESSIONS SESSIONS

- Confessions must be typed—prior to arriving at the session.
- No names will be used to identify confessions.
- Please assign a pain score from 1 to 10 on your confessions—this helps establish the tone of your confession.
- Respect all confessions.
- Respect the session—do not engage in electronic distractions, and please leave the room to answer a page.
- If you get your confession to read, do not reveal it is your confession.
- Confessions that reveal possible harm to self or others may arise. By participating in this session, we agree that the facilitator may decide to “lock down” the room for safety reasons and engage a professional therapist should the need arise.
- Facilitators of confessions agree to bring deidentified concerns to relevant faculty and/or administrators and report back on any interventions.

reflect some of the impediments to establishing accountability and accomplishing effective incident reporting. The 2012 Agency for Healthcare Research and Quality Hospital Survey on Patient Safety Culture revealed that nearly 50% of staff felt that there is a nonpunitive response to error: only 44% reported that their mistakes and event reports were not held against them, and that information on these events was not kept in their personnel file.<sup>5</sup>

With a lack of measures reflecting the incidence and quality of patient safety events, and in the face of poor reporting mechanisms, the time is ripe for adopting innovative approaches to this problem. Perhaps we need to move away from the systems approach and get back to the individual, after all.

Back at the conference table, these confessions are shared:

*Every day I am anxious, overwhelmed, and fearful about making a mistake or looking stupid. Some days it's bad; other days it's worse.*

*I mistakenly pushed an entire syringe of ephedrine thinking it was my flush syringe I usually keep on the line. The label was facing away from me, and my attending had put it on the line while I was out on break. I almost had a heart attack, but the patient barely reacted to it. I did not tell my attending.*

*A mediocre batter in the MLB probably has a higher batting average than my intubation success rate.*

*A patient yesterday asked me how long I'd been doing this, and when I said this was my first month, he got very anxious and upset.*

The more statements recited, the clearer it became that the newly minted Confessions program had far-reaching implications for the clinical learning environment. Quality improvement, on-boarding management, patient safety concerns, and a host of other issues arose from the first session. For example, the wrong drug (ephedrine) injected statement uncovers a near-miss that would otherwise not have been reported.

We started the Confessions program in 2010. In prior years, we tried other ways to understand resident stressors, including group discussions, to ease anxiety associated with the transition from medical school to residency. Michael Balint had the same thought in the 1950s when he proposed facilitated, small group discussions of the physician-patient relationship. There were ground rules: no notes were taken and all discussions were confidential. Balint's wife, Enid, a social worker and cofacilitator of these groups, commented, "... the essence ... has always been to share experiences and enable people to observe and rethink aspects of their relationships with patients and their work as doctors."

Early references to confessions exist in the Bible (Numbers 5:7). To evoke the term "confessions" seems to imply a heightened honesty and transparency to the shared information. The anonymity of confession statements adds to the residents' willingness to contribute candid vignettes. Over the years, Confessions sessions have succeeded when (1) sessions were led by experienced facilitators; (2) a safe environment was ensured for the sessions; (3) ground rules were clearly defined and not violated; and (4) statements that warranted intervention resulted in clinical learning environment improvements, and quality and patient safety improvements, with measurable outcomes that were reported back to the participants. See the BOX for our rules for the sessions.

Our experience has provided us with some specific "dos and don'ts." We have found that the ideal group consists of 8 to 12 participants, and the optimal setting allows for everyone to sit in a circle. Allotting about 1 hour for each session seems to be enough time, and we try our best to get through everyone's confessions. The facilitator should try not to hijack the conversation and is advised to act more as a guide and to not be overly authoritarian. The facilitator also should take notes to ensure that necessary and timely follow-up occurs. For example, a confession about a drug error stimulated a conversation among the

residents about how drugs are located in the storage cart. Rather than blame the confessor about the drug error, a systemic problem was sought and a recommendation was made to reorganize the drug drawers. The facilitator followed up with the pharmacist and then reported back to the group that the desired change had occurred.

We also have met and overcome some challenges in holding these sessions. For example, in one program, a designated facilitator jeopardized the group trust when he stated, “You guys are in residency; it’s supposed to be challenging. Stop whining about it.” He then proceeded to exit the room. The residents felt betrayed, stating they were encouraged to confess their fears, and were assured that there would be no negative responses. This experience threatened to derail the safety and trust in the environment. In another program, when difficulty initiating conversations was noted, a facilitator slipped a “fake confession” into the pool. The planted confession was regarding the false documentation of heart and lung auscultation when it had not actually been performed. This particular issue had arisen with a previous group of residents, so the program director believed it would help stimulate frank dialogue—which it did.

Our use of Confessions has provided an opportunity to reflect, discuss, and admit without fear of punitive actions. It allows for early intervention on the issues that are relevant to physician trainees. If used effectively by program directors, and by department chairs and division leaders with faculty, the Confessions program could guide modification of patient care and further quality care efforts in focused

and specific ways that are currently not being addressed in any other manner.

## References

1. Reznick MA, Barton BA. Improved incident reporting following the implementation of a standardized emergency department peer review process. *Int J Qual Health Care*. 2014;26(3):278–286.
2. Rozich JD, Haraden CR, Resar RK. Adverse drug event trigger tool: a practical methodology for measuring medication related harm. *Qual Saf Health Care*. 2003;12(3):194–200.
3. Leape LL, Berwick DM. Five years after *To Err Is Human*: what have we learned? *JAMA*. 2005;293(19):2384–2390.
4. Rowin EJ, Lucier D, Pauker SG, Kumar S, Chen J, Salem DN. Does error and adverse event reporting by physicians and nurses differ? *Jt Comm J Qual Patient Saf*. 2008;34(9):537–545.



**Suzanne B. Karan, MD**, is Associate Professor of Anesthesiology, Department of Anesthesiology, University of Rochester School of Medicine & Dentistry; **Jeffrey S. Berger, MD, MBA**, is Associate Professor of Anesthesiology, Department of Anesthesiology & Critical Care Medicine, George Washington University School of Medicine & Health Sciences; and **Michael Wajda, MD**, is Associate Professor of Anesthesiology, Department of Anesthesiology, New York University Langone Medical Center.

Corresponding author: Suzanne B. Karan, MD, University of Rochester School of Medicine & Dentistry, Department of Anesthesiology, Box 604, 601 Elmwood Avenue, Rochester, NY 14642, 585.275.1384, fax 585.276.0122, [suzanne\\_karan@urmc.rochester.edu](mailto:suzanne_karan@urmc.rochester.edu)

**Queries for jgme-07-04-02**

- 1. Author: This article has been edited for grammar, style, and usage. Please compare it with your original document and make corrections on these pages. Please limit your corrections to substantive changes that affect meaning. If no change is required in response to a question, please write “OK as set” in the margin. Copy editor**
- 2. Author: In sentence “A few programs have been successful...” what is “more is clearly needed” referring to? Research? Programs? Copy editor**