



ORAL & MAXILLOFACIAL SURGERY

POSTOPERATIVE INSTRUCTIONS FOLLOWING DENTAL IMPLANTS or RELATED SURGERY

- 1. First 30 minutes after procedure:** If asked to do so, Bite on gauze for 30 minutes after your procedure, to help stop bleeding. It is best that you do this without checking the site until the 30 minutes is up. Do not chew on the gauze, as continued pressure is desired. During this time, do not speak or do other activities that could dislodge the gauze packs. If bleeding continues after 30 minutes, bite on fresh gauze as follows: dampen the gauze first so that it is moist (not wet) and fold it twice. Place the dampened gauze on the extraction sites for another 30 minutes. Repeat as necessary. A small amount of continued blood, which tinges the saliva, is expected during the first 24 hours and does not require further pressure. If bleeding restarts after it has stopped, use gauze pressure again or consider gently biting on a dry tea bag placed over the extraction site for 30 minutes. If the bleeding does not stop despite repeated attempts as above, please call 275-5531.
- 2. Avoid excessive exertion** for 72 hours after your procedure. No bending, lifting, strenuous activity, sports or gym-class. After this time, resume normal activity gradually as you become comfortable.
- 3. Mouth opening will be limited** during the first week to ten days after surgery. This is expected due to swelling and tightness of the jaw muscles.
- 4. Bruising** (black and blue areas) may develop after surgery. Should this occur, expect bruising to resolve after the first week to ten days.
- 5. Earache or TMJ (jaw joint) tenderness** may develop that will gradually resolve over the first week.
- 6. NO ORAL CARE** during the first 24 hours. Beginning on the day after surgery, rinse your mouth after meals (or 4-5 times per day) with warm salt water (1/2 teaspoon of salt in an 8oz. glass of warm water). A prescription mouth rinse may be given in selected cases. Begin tooth brushing on the day after surgery, as you normally would however this should be performed gently in the area of extractions.
- 7. DO NOT USE STRAWS** for the first week after surgery. Straws create suction pressure inside your mouth and could dislodge the blood clots that normally form in the extraction sites.
- 8. DO NOT SMOKE** during the first 7 days after surgery. Smoking greatly increases the risk of dry socket and other healing complications.
- 9. SWELLING** is normal and will maximize 48 – 72 hours after surgery. Apply ice in a damp towel against your cheeks for the first 24-48 hours, as follows: 20 minutes on 40 minutes off, then repeat. Keep your head elevated with 2 or 3 pillows when lying down during the first 24 hours. This will reduce bleeding, swelling and improve your level of comfort. May apply Vaseline^R or Chapstick^R to the lips to keep them moist.
- 10. Mild to moderate pain** may be encountered after the local anesthetic wears off. If not contraindicated by your health history, take two Extra-Strength Tylenol^R (500mg ea.) or two Motrin^R/Advil^R/Ibuprofen (200mg ea.). Avoid aspirin-containing products since these may lead to more bleeding. For moderate to severe pain, take the prescription medication as prescribed for you. **Do not operate heavy machinery or make important decisions while on narcotics.**
- 11.** In case of uncontrolled bleeding, extreme pain or unusual circumstances, please call the office immediately.



POSTOPERATIVE DIET SUGGESTIONS

1. **Avoid eating or drinking anything within the first hour after surgery.** After this time, drink lots of liquids and eat soft, cool foods. Avoid excessively hot liquids and foods for the first 24 hours.
2. During the first two weeks, **avoid hard, crispy foods such as chips, hard pretzels, and popcorn.**
3. The following food choices are recommended (“liquid foods on the day of the procedure, soft “mushy” foods during the first week advancing to regular food as tolerated”):
 - a. **During the first two weeks**, blenderized foods, soup broth, well-cooked pasta, eggs, yogurt, cottage cheese, Jell-O, pudding, custard, applesauce, bananas and ice cream.
 - b. **After the first two weeks**, gradually increase your diet to include: fish, chicken, meatloaf, mashed potatoes, cooked vegetables, soups and cereals, etc... and then to regular food as tolerated.

ORAL & MAXILLOFACIAL SURGERY

SINUS PRECAUTIONS

1. Keep your head elevated during sleep with two to three pillows.
2. **Do NOT blow your nose**
3. **If you sneeze** do not sneeze through the nose and do not “hold back.” You may sneeze with your mouth open.
4. Expect slight nose bleeds. Call the office immediately for heavy bleeding from the nose that you are unable to control with pressure.
5. **Do NOT** do anything that requires a sucking-in or blowing-out motion such as smoking, use of straws or blowing up balloons.
6. Take all prescribed and/or recommended medications as directed.